

- 关产品安全所. 第三次中国城市儿童哮喘流行病学调查[J]. 中华儿科杂志, 2013, 51(10): 729-735.
- [2] Zholos AV. TRP channels in respiratory pathophysiology: the role of oxidative, chemical irritant and temperature stimuli[J]. Curr Neuropharmacol, 2015, 13(2): 279-291.
- [3] Chen CL, Li H, Xing XH, et al. Effect of TRPV1 gene mutation on bronchial asthma in children before and after treatment[J]. Allergy Asthma Proc, 2015, 36(2): e29-e36.
- [4] Locke NR, Royce SG, Wainwright JS, et al. Comparison of airway remodeling in acute, subacute, and chronic models of allergic airways disease[J]. Am J Respir Cell Mol Biol, 2007, 36(5): 625-632.
- [5] Delescluse I, Mace H, Adcock JJ. Inhibition of airway hyper-responsiveness by TRPV1 antagonists (SB-705498 and PF-04065463) in the unanaesthetized ovalbumin-sensitized guinea pig[J]. Br J Pharmacol, 2012, 166(6): 1822-1832.
- [6] Cantero-Recasens G, Gonzalez JR, Fandos C, et al. Loss of function of transient receptor potential vanilloid 1 (TRPV1) genetic variant is associated with lower risk of active childhood asthma[J]. J Biol Chem, 2010, 285(36): 27532-27535.
- [7] Ma J, Altomare A, Guarino M, et al. HCl-induced and ATP-dependent upregulation of TRPV1 receptor expression and cytokine production by human esophageal epithelial cells[J]. Am J Physiol Gastrointest Liver Physiol, 2012, 303(5): G635-G645.
- [8] Wang Z, Yang Y, Yang H, et al. NF- κ B feedback control of JNK1 activation modulates TRPV1-induced increases in IL-6 and IL-8 release by human corneal epithelial cells[J]. Mol Vis, 2011, 17: 3137-3146.
- [9] Tsiligianni Z, Hillas G, Bakakos P, et al. Sputum interleukin-13 as a biomarker for the evaluation of asthma control[J]. Clin Exp Allergy, 2016, 46(7): 923-931.
- [10] Dunn R, Szeffler SJ. Severe asthma in pediatric patients. Pathophysiology and unmet needs[J]. Ann Am Thorac Soc, 2016, 13 Suppl 1: S103-S104.
- (本文编辑: 万静)

· 消息 ·

本刊关于实验动物样本数的规定

为了提高杂志的学术质量,《中国当代儿科杂志》决定从 2016 年 8 月起,对凡是在本刊刊出的动物实验研究论文中实验动物样本数做出如下规定,请广大作者注意并严格执行。对于未达到以下要求的动物实验性文章投稿,本刊则一律不再接收。

(1) 活体动物实验每组动物数不得少于 8 只;

(2) 对于细胞或分子水平检测方法,可适当放宽对样本数的要求,如:PCR 法检测 mRNA 水平,每个单元组的样本数不得少于 6 例;Western blot 法检测蛋白水平,每个单元组的样本数不得少于 3 例;

(3) 组织切片,要求每个样本组织的切片数不得少于 3 张,每张切片随机观察视野不得少于 5 个。

《中国当代儿科杂志》编辑部
2016 年 8 月