Child Neurol, 2014, 29(2): 265-268.
[13] Swoboda KJ, Kanavakis E, Xaidara A, et al. Alternating hemiplegia of childhood or familial hemiplegic migraine? A novel ATP1A2 mutation[J]. Ann Neurol, 2004, 55(6): 884-887.
[14] de Vries B, Stam AH, Beker F, et al. CACNA1A mutation linking hemiplegic migraine and alternating hemiplegia of childhood[J]. Cephalalgia, 2008, 28(8): 887-891.
[15] Vila-Pueyo M, Pons R, Raspall-Chaure M, et al. Clinical and genetic analysis in alternating hemiplegia of childhood: ten new patients from Southern Europe[J]. J Neurol Sci, 2014, 344(1-2): 37-42.
[16] Weller CM, Leen WG, Neville BG, et al. A novel SLC2A1 mutation linking hemiplegic migraine with alternating hemiplegia of childhood[J]. Cephalalgia, 2015, 35(1): 10-15.
[17] Vuillaumier-Barrot S, Panagiotakaki E, Le Bizec C, et al. Absence of mutation in the SLC2A1 gene in a cohort of patients with alternating hemiplegia of childhood (AHC)[J]. Neuropediatrics, 2010, 41(6): 267-269.
[18] De Grandis E, Stagnaro M, Biancheri R, et al. Lack of SLC2A1 (glucose transporter 1) mutations in 30 Italian patients with alternating hemiplegia of childhood[J]. J Child Neurol, 2013, 28(7): 863-866.
[19] de Vries B, Haan J, Stam AH, et al. Alternating hemiplegia of childhood: no mutations in the glutamate transporter EAAT1[J]. Neuropediatrics, 2006, 37(5): 302-304.
[20] Heinzen EL, Swoboda KJ, Hitomi Y, et al. De novo mutations in ATP1A3 cause alternating hemiplegia of childhood[J]. Nat Genet, 2012, 44(9): 1030-1034.
[21] Rosewich H, Thiele H, Ohlenbusch A, et al. Heterozygous de-novo mutations in ATP1A3 in patients with alternating hemiplegia of childhood: a whole-exome sequencing gene-identification study[J]. Lancet Neurol, 2012, 11(9): 764-773.
[22] Dobretsov M, Stimers JR. Neuronal function and alpha3 isoform of the Na/K-ATPase[J]. Front Biosci, 2005, 10: 2373-2396.
[23] Panagiotakaki E, De Grandis E, Stagnaro M, et al. Clinical profile of patients with ATP1A3 mutations in Alternating Hemiplegia of Childhood-a study of 155 patients[J]. Orphanet J Rare Dis, 2015, 10: 123.
[24] Ishii A, Saito Y, Mitsui J, et al. Identification of ATP1A3 mutations by exome sequencing as the cause of alternating hemiplegia of childhood in Japanese patients[J]. PLoS One, 2013, 8(2): e56120.

（本文编辑：俞燕）

·消息·

《实用儿童脑病学》新书介绍

儿童脑损伤和脑病是危害儿童健康成长和人口素质的主要疾病，严重影响儿童社会交往能力、适应能力、注意力和学习能力，甚至导致癫痫、视听障碍、语言障碍、智力障碍、脑性瘫痪、植物状态等神经伤残。

由陈光福教授主编、人民卫生出版社出版的《实用儿童脑病学》，全书共 30 章，139 万字，460 多幅图（部分彩图），涉及儿童神经发育、神经解剖、神经影像学检查、脑功能检查、神经学检查评估、神经保护治疗、神经免疫治疗、神经修复治疗与康复治疗，以及各种儿童脑疾病的病因、发病机制、病理、临床表现、诊断方法与治疗进展等内容。

本书编委为从事儿童神经发育基础、神经影像、神经电生理、神经内科、神经外科、儿童康复和新生儿等专业的专家，专家们历时 2 年精心撰写，反复修改、补充，力求反映本领域近 5 年国内外的最新研究进展、诊治指南和专家共识，以及作者的临床经验与研究成果，并对新的探索性治疗应用前景进行了述评和展望。全书图文并茂、条理清晰、内容丰富、可读性强，是一部具有先进性、科学性和实用性的专业参考书。

有意购买者请联系王波医师，电话：13823108103，邮箱：66286432@qq.com。