

- analysis of umbilical blood samples in 2036 newborns[J]. Int Reprod Health Fam Plan, 2014, 33: 478-480.
- [7] Zeitlin J, Mohangoo A, Cuttini M, et al. The European perinatal health report: comparing the health and care of pregnant women and newborn babies in Europe[J]. J Epidemiol Community Health, 2009, 63(9): 681-682.
- [8] Abu-Amero KK, Hellani AM, Salih MA, et al. A de novo marker chromosome derived from 9p in a patient with 9p partial duplication syndrome and autism features: genotype-phenotype correlation[J]. BMC Med Genet, 2010, 11: 135.
- [9] Amasdl S, Natiq A, Elalaoui SC, et al. Insulin-like growth factor type 1 deficiency in a Moroccan patient with de novo inverted duplication 9p24p12 and developmental delay: a case report[J]. J Med Case Rep, 2016, 10(1): 122.
- [10] Hou QF, Wu D, Chu Y, et al. Clinical findings and molecular cytogenetic study of de novo pure chromosome 9p deletion: Pre and postnatal diagnosis[J]. Taiwan J Obstet Gynecol, 2016, 55(6): 867-870.
- [11] Onesimo R, Orteschi D, Scalzone M, et al. Chromosome 9p deletion syndrome and sex reversal: novel findings and redefinition of the critically deleted regions[J]. Am J Med Genet A, 2012, 158A(9): 2266-2271.
- [12] Vialard F, Ottolenghi C, Gonzales M, et al. Deletion of 9p associated with gonadal dysfunction in 46,XY but not in 46,XX human fetuses[J]. J Med Genet, 2002, 39(7): 514-518.
- [13] Witters I, Vermeesch JR, Moerman PH, et al. Partial trisomy 3p/monosomy 9p with sex reversal[J]. Ultrasound Obstet Gynecol, 2004, 23(4): 418-419.
- [14] Chen CP, Lin CL, Chen LL, et al. Prenatal diagnosis of mosaic ringchromosome 9[J]. Prenat Diagn, 2006, 26(9): 870-871.
- [15] Hou WC, Chen CP, Hwang KS, et al. Prenatal diagnosis of a de novo 9p terminal chromosomal deletion in a fetus with major congenital anomalies[J]. Taiwan J Obstet Gynecol, 2014, 53(4): 602-605.
- [16] 杨尧, 王芳, 王春枝, 等. 孤独症患儿1例遗传学分析[J]. 中国儿童保健杂志, 2012, 20(10): 893-895.
- (本文编辑: 俞燕)

· 消息 ·

本刊关于实验动物样本数的规定

为了提高杂志的学术质量,《中国当代儿科杂志》决定从2016年8月起,对凡是在本刊刊出的动物实验研究论文中实验动物样本数做出如下规定,请广大作者注意并严格执行。对于未达到以下要求的动物实验性文章投稿,本刊则一律不再接收。

(1) 活体动物实验每组动物数不得少于8只;

(2) 对于细胞或分子水平检测方法,可适当放宽对样本数的要求,如:PCR法检测mRNA水平,每个单元组的样本数不得少于6例;Western blot法检测蛋白水平,每个单元组的样本数不得少于3例;

(3) 组织切片,要求每个样本组织的切片数不得少于3张,每张切片随机观察视野不得少于5个。

《中国当代儿科杂志》编辑部
2018年1月